

Five Things Physicians and Patients Should Question

1

Do not order a screening hip ultrasound to rule out developmental hip dysplasia or developmental hip dislocation if the baby has no risk factors and has a clinically stable hip examination.

Hip dysplasia/dislocation is relatively rare, with incidence of approximately 7 per 1,000 births. Studies have shown that universal screening programs for developmental hip instability using ultrasounds to assess otherwise normal appearing hips have a nearly negligible positive yield. There is a substantial false positive rate, with an associated increase in treatment rate, suggesting that babies without hip pathology are being treated. When there are no physical findings or underlying risk factors for hip dysplasia/dislocation in a newborn, a hip ultrasound is costly, time-intensive and the findings may be misleading to parents and physicians. This recommendation is in accordance with the 2016 AAP clinical report regarding the use of ultrasound in early detection of developmental dysplasia of the hip (see reference: "Evaluation and Referral for Developmental Dysplasia of the Hips in Infants").

2

Do not order radiographs or advise bracing or surgery for a child less than 8 years of age with simple in-toeing gait.

Mild in-toeing is usually a physiologic phenomenon reflecting ongoing maturation of the skeleton. Metatarsus adductus, femoral anteversion, and tibial torsion all contribute to in-toeing and tend to improve with growth. Simply monitoring gait for continued improvement at normal well child examination intervals is adequate until the age of 7–8 unless there is severe tripping and falling or asymmetry. It is not possible to alter the natural evolution using physical therapy, bracing or shoe inserts.

3

Do not order custom orthotics or shoe inserts for a child with minimally symptomatic or asymptomatic flat feet.

Flexible flat feet are normal physiologic variants commonly found in children and adults. Unlike a painful or rigid flatfoot that requires further workup, if an arch is present when standing on tiptoe, the foot can be managed with observation or over-the-counter orthotics. The use of custom orthotic devices to provide support for the foot does not aid in the development of the arch.

4

Do not order advanced imaging studies (MRI or CT) for most musculoskeletal conditions in a child until all appropriate clinical, laboratory and plain radiographic examinations have been completed.

History, physical examination, and appropriate radiographs remain the primary diagnostic modalities in pediatric orthopaedics, as they are both diagnostic and prognostic for the great majority of pediatric musculoskeletal conditions. Examples of such conditions would include, but not be limited to, the work up of injury or pain (spine, knees and ankles), possible infection, and deformity. MRI examinations and other advanced imaging studies are costly, frequently require sedation in the young child (5 years old or less), and may not result in appropriate interpretation if clinical correlations cannot be made. Many conditions require specific MRI sequences or protocols best ordered by the specialist who will be treating the patient. Inappropriately obtained MRIs may need to be repeated in those circumstances. Additionally, a significant dose of radiation is delivered to the patient during a CT scan, so their utility in a specific case would be best confirmed prior to ordering. Therefore, in those conditions where advanced imaging is indicated, it has greater value when it is used to answer a specific question that arises from a thorough clinical and appropriate radiographic evaluation. Additionally, if you believe findings warrant additional advanced imaging, discuss with the consulting orthopaedic surgeon to make sure the optimal studies are ordered.

5

Do not order follow-up X-rays for buckle (or torus) fractures if they are no longer tender or painful.

Buckle (torus) fractures are very common injuries in young children, especially in the distal radius. The fracture is one of compression, where the metaphyseal bone impacts on itself, and actually becomes denser. These fractures are inherently stable and do not necessarily require a formal cast, unless severe pain or fracture instability necessitates a cast for 4 weeks. Instead immobilization with a simple wrist brace or removable splint is often preferable. The mild cortical angular deformity reliably remodels over time and requires no intervention or monitoring. If the fracture is non-tender to palpation at 4 weeks post-injury, no follow-up radiograph is required, and full activities may be resumed.

How This List Was Created

The Pediatric Orthopaedic Society of North America (POSNA) Evidence Based Medicine Committee and the Advocacy Committee worked together during 2014 and 2015 to develop five items in the practice of Pediatric Orthopaedics of tests or procedures that should not be done routinely. Approximately 20 members of these two committees participated in the process. Each surgeon, in a blinded fashion, submitted 5 items each from their practices and experience of tests or procedures that they found were commonly over-utilized. The items were tallied in order of number of times that item was listed by each surgeon. A total of 30 items were submitted. Both committees then agreed on final list of 5 items based of frequency of responses and importance of the condition. The Evidence Based Committee reviewed the appropriate literature to provide references and support for each item. The Executive Committee of the Orthopaedic Section of the American Academy of Pediatrics (AAP) reviewed the 5 listed items and provided further feedback. POSNA Board of Directors provided further feedback and voted on the final list. Various expert committees and sections of the AAP reviewed and approved the list. The AAP Executive Committee granted final approval of the list.

Sources

- 1 Mahan ST, Katz JN, Kim YJ. To screen or not to screen? A decision analysis of the utility of screening for developmental dysplasia of the hip. *J Bone Joint Surg Am.* 2009 Jul;91(7):1705-1719.
- 2 Laborie LB, Markestad TH, Davidsen H, Bruras KR, Aukland SM, Bjortlykke JA, Reigstad H, Indrekvam K, Lehmann TG, Engesaeter IO, Engesaeter LB, Rosendahl K. Selective ultrasound screening for developmental hip dysplasia: effect on management and late detected cases. A prospective study during 1991-2006. *Pediatr Radiol.* 2014 Apr;44 (4): 410-424.
- 3 Shorter D, Hong T, Osborn DA. Cochrane Review: Screening programs for developmental dysplasia of the hip in newborn infants. *Evid Based Child Health.* 2013; 8(1): 11-54.
- 4 Shaw BA, Segal LS. Section on Orthopaedics. Evaluation and referral for developmental dysplasia of the hip in infants. *Pediatrics* 2016; 138(6).
- 5 Fabry G, Cheng LX, Molenaers G. Normal and abnormal torsional development in children. *Clinical Orthopaedics and Related Research.* May 1994; (301):22-26.
- Fabry G, MacEwen GD, Sharnds AR, Jr. Torsion of the femur: A follow up study in normal and abnormal conditions. *J Bone Joint Surg. Am.* Dec 1973;55(8):1726-1738.
- Lincoln TL, Suen PW. Common rotational variations in children. The Journal of the American Academy of Orthopaedic Surgeons. Sep-Oct 2003; 11(5):312-320.
- Staheli LT, Corbett M, Wyss C, King H. Lower-extremity rotational problems in children. Normal values to guide management. *J Bone Joint Surg Am.* Jan 1985;67(1):39-47.
- Svenningsen S, Apalset K, Terjesen T, Anda S. Regression of femoral anteversion. A prospective study of in-toeing of children. *Acta Orthopaedica Scandinavica.* Apr 1989;60(2):170-173.
- Wenger DR, Mauldin D, Speck G, Morgan D, Lieber RL. Corrective shoes and inserts as treatment for flexible flatfoot in infants and children. *J Bone Joint Surg Am.* 1989 Jul;71(6):800-810.
- Staheli LT, Chew DE, Corbett M. The longitudinal arch: A survey of eight hundred and eighty-two feet in normal children and adults. *J Bone Joint Surg Am.* 1987 Mar;69(3):426-428.
- Piccolo CL, Galluzzo M, Ianniello S, Trinci M, Russo A, Rossi E, Zeconlini M, Laporta A, Guglielmi G, Muielo V. Pediatric musculoskeletal injuries: role of ultrasound and magnetic resonance imaging. *Musculoskelet Surg.* 2017 Mar; 101(Suppl 1):85-102.
- LaBella CR, Henrikus W, Hewett TE. Anterior cruciate ligament Injuries: Diagnosis, Treatment, and Prevention. *Pediatrics* 2014;133(5):e1437-e1450.
- Tuite MJ, Kransdorf MJ, Beaman FD, Adler RS, Amini B, Appel M, Bernard SA, Dempsey ME, Fries IB, Greenspan BS, Khurana B, Mosher TJ, Walker EA, Ward RJ, Wessell DE, Weissman BN. ACR Appropriateness Criteria[®] Acute Trauma to the Knee. Available at <https://acsearch.acr.org/docs/69419/Narrative/> American College of Radiology. Revised 2014.
- Deyle GD. The role of MRI in musculoskeletal practice: a clinical perspective. *J Man Manip Ther.* 2011 Aug;19(3):152-161
- Bateni C, Bindra J, Haus B. MRI of sports injuries in children and adolescents: what's different from adults. *Current Radiology Reports.* 2014;2:45.
- Symons S, Rowsell M, Bhowal B, Diass JJ. Hospital versus home management of children with buckle fractures of the distal radius: A prospective randomized trial. *J Bone Joint Surg Br.* 2001;83:556-560.
- Van Bosse HJ, Patel RJ, Thacker M, Sala DA. Minimalistic approach to treating wrist torus fractures. *J Pediatric Orthop.* 2005 Jul-Aug;25(4):495-500.
- Williams KG, Smith G, Luhmann SJ, Mao J, Gunn JD, Luhmann JD. A randomized controlled trial of cast versus splint for distal radial buckle fracture: An evaluation of satisfaction, convenience, and preference. *Pediatric Emergency Care.* 2013 May;29(5):555-559.

About the ABIM Foundation

The mission of the ABIM Foundation is to advance medical professionalism to improve the health care system. We achieve this by collaborating with physicians and physician leaders, medical trainees, health care delivery systems, payers, policymakers, consumer organizations and patients to foster a shared understanding of professionalism and how they can adopt the tenets of professionalism in practice.



To learn more about the ABIM Foundation, visit www.abimfoundation.org.

About the AAP-Section on Orthopaedics

The American Academy of Pediatrics is an organization of 66,000 primary care pediatricians, pediatric medical subspecialists and pediatric surgical specialists dedicated to the health, safety and well-being of infants, children, adolescents and young adults. The Section on Orthopaedics was founded over 40 years ago for the primary purpose of improving the musculoskeletal health of children through mentorship, education, research, advocacy, and service. The Section includes over 150 pediatric orthopaedic surgeons and sports medicine physicians who often collaborate with members of international societies such as the Pediatric Orthopaedic Society of North America.



About the Pediatric Orthopaedic Society of North America

The Pediatric Orthopaedic Society of North America (POSNA) is a not-for-profit professional organization of over 1,400 surgeons, physicians, and allied health members passionately dedicated to advancing musculoskeletal care for children and adolescents through education, research, quality, safety and value initiatives, advocacy, and global outreach to children in underserved areas.



For more information or to see other lists of Things Providers and Patients Should Question, visit www.choosingwisely.org.