

Five Things Physicians and Patients Should Question

1

Avoid routine spinal imaging in the absence of clear clinical indicators for patients with acute low back pain of less than six (6) weeks duration.

Multidisciplinary evidence-based guidelines recommend against the routine use of spinal imaging for patients with acute low back pain of less than six weeks duration in the absence of clear clinical indicators. Such indicators include, but are not limited to, history of cancer, fracture or suspected fracture based on clinical history, progressive neurologic symptoms, and infection. Doctors of chiropractic must also consider conditions that potentially preclude a dynamic thrust to the spine, which include but are not limited to, osteopenia, osteoporosis, axial spondyloarthritis and tumors. Unnecessary imaging incurs monetary cost, exposes the patient to ionizing radiation, and can result in labeling patients with conditions that are not clinically meaningful, creating a false sense of vulnerability and disability. Indeed, several studies have shown that the routine use of radiographs in the care of low back pain may result in worse outcomes than without their use.

2

Do not perform repeat imaging to monitor patients' progress.

With few exceptions (e.g., the long-term management of idiopathic scoliosis) radiographic findings should not be used as outcome measures for low-back pain. There is currently no data available to support a relationship between changes in alignment or other structural characteristics and patient improvement. This practice increases costs, exposes patients unnecessarily to ionizing radiation and may distract from more meaningful outcomes. Furthermore, there is no known correlation between performing routine or repeat imaging studies to monitor a patient's condition and improved clinical outcomes or meaningful changes in patient management. Repeat imaging is appropriate only if strong clinical indications exist, such as a major change in diagnosis, documented worsening of symptoms or significant progression of disease. Failure to respond to treatment is not an indication for repeat imaging.

3

Avoid protracted use of passive or palliative physical therapeutic modalities for low-back pain disorders unless they support the goal(s) of an active treatment plan.

Passive physical therapeutic modalities are defined as those interventions applied to a patient with no active participation on the part of the patient. These include heat, cold, electrical stimulation and ultrasound. These passive therapies can play an important role in facilitating patient participation in an active treatment program. However, the use of passive therapies untethered to the goal of increasing physical activity can be harmful, as it can lead to patient inactivity, prolonged recovery and increased costs. For any patient with a low-back pain disorder to achieve an optimal clinical outcome, an essential element is to restore, maintain or increase the level of physical activity. The evidence demonstrates that both general physical activity (e.g., walking, jogging, biking) and specific exercise regimens are effective in treating and preventing low-back pain and may lead to better outcomes when combined with spinal manipulation.

4

Do not provide long-term pain management without a psychosocial screening or assessment.

There is a high probability that any person with a chronic pain syndrome has a concomitant psychological disorder, most notably depression and/or anxiety. The relationship between chronic pain and depression/anxiety is well established. The causal arrow between pain and these disorders can point in either direction and over time may form a positive feedback loop between these two elements. Screening tools are available that will aid in the detection of potential depression/anxiety, and, when indicated, a referral may be most appropriate for more extensive evaluation and treatment. In addition, lesser psychological factors such as catastrophizing and fear avoidance behavior may interfere with a patient's recovery and should be recognized by the clinician. Recognizing indicators of patient psychosocial health behavioral factors can affect a patient's recovery and/or compliance with treatment and may decrease the risk of developing chronic illness/pain. Tools such as StarTBack 9 screening tool, PHQ-9 depression scale and the Fear Avoidance Belief Questionnaire are examples.

5

Do not prescribe lumbar supports or braces for the long-term treatment or prevention of low-back pain.

While there may be limited benefit in the short term, the prolonged use of lumbar supports is not supported by the literature for the treatment or prevention of low-back pain. Numerous systematic reviews have found limited to no value for their use in this context. The literature clearly demonstrates that such passive therapies are contrary to the currently accepted central principle of low-back pain care, which is that the patient must engage in an active rehabilitative regimen to achieve the best outcomes.

How This List Was Created

The American Chiropractic Association (ACA) utilized its Committee on Quality Assurance and Accountability (CQAA) to serve as an expert task force of doctors of chiropractic (DCs) to identify areas/items common to the practice of chiropractic for which recommendations were supported by clinical research and would result in high-value, cost-effective services and improved patient outcomes. A literature search was conducted and the task force collaboratively identified a draft list of six recommendations based upon established *Choosing Wisely*[®] criteria. The list was submitted to the ACA Board of Governors for initial review. After further refinement, the final list of five strategies was selected, submitted to and approved by the ACA Board of Governors.

ACA's disclosure and conflict of interest policy can be found at www.acatoday.org.

Sources

1

Chou R, Fu R, Carrino JA, Deyo RA. Imaging strategies for low-back pain: systematic review and meta-analysis. *Lancet*. 2009 Feb 7;373(9662):463-72.

Bussi eres AE, Taylor JA, Peterson C. Diagnostic imaging practice guidelines for musculoskeletal complaints in adults—an evidence-based approach—part 3: spinal disorders. *J Manipulative Physiol Ther*. 2008 Jan;31(1):33-88.

Kendrick D, Fielding K, Bentley E, Miller P, Kerslake R, Pringle M. The role of radiography in primary care patients with low back pain of at least 6 weeks duration: a randomised (unblinded) controlled trial. *Health Technol Assess*. 2001;5(30):1-69.

National Guideline Clearinghouse (NGC). Guideline summary: ACR Appropriateness Criteria[®] low back pain. In: National Guideline Clearinghouse (NGC) [Web site]. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); [2016 Jan 22]. Available from: <https://www.acr.org/Clinical-Resources/ACR-Appropriateness-Criteria>

Brinjikji W, Luetmer PH, Comstock B, Bresnahan BW, Chen LE, Deyo RA, Halabi S, Turner JA, Avins AL, James K, Wald JT, Kallmes DF, Jarvik JG. Systematic literature review of imaging features of spinal degeneration in asymptomatic populations. *AJNR Am J Neuroradiol*. 2015 Apr;36(4):811-6.

2

Brinjikji W, Luetmer PH, Comstock B, Bresnahan BW, Chen LE, Deyo RA, Halabi S, Turner JA, Avins AL, James K, Wald JT, Kallmes DF, Jarvik JG. Systematic literature review of imaging features of spinal degeneration in asymptomatic populations. *AJNR Am J Neuroradiol*. 2015 Apr;36(4):811-6.

Matsumoto M, Okada E, Toyama Y, Fujiwara H, Momoshima S, Takahata T. Tandem age-related lumbar and cervical intervertebral disc changes in asymptomatic subjects. *Eur Spine J*. 2013 Apr;22(4):708-13.

Okada E, Matsumoto M, Fujiwara H, Toyama Y. Disc degeneration of cervical spine on MRI in patients with lumbar disc herniation: comparison study with asymptomatic volunteers. *Eur Spine J*. 2011 Apr;20(4):585-91.

Chou R, Fu R, Carrino JA, Deyo RA. Imaging strategies for low-back pain: systematic review and meta-analysis. *Lancet*. 2009 Feb 7;373(9662):463-72.

Kendrick D, Fielding K, Bentley E, Kerslake R, Miller P, Pringle M. Radiography of the lumbar spine in primary care patients with low back pain: randomised controlled trial. *BMJ* 2001 Feb 17; 322(7283): 400-5.

Bussi eres AE, Taylor JA, Peterson C. Diagnostic imaging practice guidelines for musculoskeletal complaints in adults-an evidence-based approach-part 3: spinal disorders. *J Manipulative Physiol Ther*. 2008 Jan;31(1):33-88.

National Guideline Clearinghouse (NGC). Guideline summary: ACR Appropriateness Criteria[®] low back pain. In: National Guideline Clearinghouse (NGC) [Web site]. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); [2016 Jan 22]. Available from: <https://www.acr.org/Clinical-Resources/ACR-Appropriateness-Criteria>

3

Ebadi S, Henschke N, Nakhostin Ansari N, Fallah E, van Tulder MW. Therapeutic ultrasound for chronic low-back pain. *Cochrane Database Syst Rev*. 2014 Mar 14;(3):CD009169.

McGregor AH, Probyn K, Cro S, Dor e CJ, Burton AK, Balagu e F, Pincus T, Fairbank J. Rehabilitation following surgery for lumbar spinal stenosis. *Cochrane Database Syst Rev*. 2013 Dec;(12):CD009644.

Khadiilkar A, Odebiyi DO, Brosseau L, Wells GA. Transcutaneous electrical nerve stimulation (TENS) versus placebo for chronic low-back pain. *Cochrane Database Syst Rev*. 2008 Oct 8;(4):CD003008.

Steffens D, Maher CG, Pereira LS, Stevens ML, Oliveira VC, Chapple M, Teixeira-Salmela LF, Hancock MJ. Prevention of Low Back Pain: A Systematic Review and Meta-analysis. *JAMA Intern Med*. 2016 Feb;176(2):199-208.

Chou R, Deyo R, Friedly J, et al. Noninvasive Treatments for Low Back Pain. Comparative Effectiveness Review No. 169. [Internet]. Rockville (MD): Agency for Healthcare Research and Quality (US); 2016 Feb. (Comparative Effectiveness Reviews, No. 169.) [cited 2017 May 4]. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK350276/>

Cancelliere C, Donovan J, Stochkendahl MJ, Biscardi M, Ammendolia C, Myburgh C, Cassidy JD. Factors affecting return to work after injury or illness: best evidence synthesis of systematic reviews. *Chiropr Man Therap*. 2016 Sep 8;24(1):32.

Kamper SJ, Apeldoorn AT, Chiarotto A, Smeets RJ, Ostelo RW, Guzman J, van Tulder MW. Multidisciplinary biopsychosocial rehabilitation for chronic low back pain. *Cochrane Database Syst Rev*. 2014 Sep 2;(9):CD000963.

Wertli MM, Eugster R, Held U, Steurer J, Kofmehl R, Weiser S. Catastrophizing-a prognostic factor for outcome in patients with low back pain: a systematic review. *Spine J*. 2014 Nov 1;14(11):2639-57.

Taylor JB, Goode AP, George SZ, Cook CE. Incidence and risk factors for first-time incident low back pain: a systematic review and meta-analysis. *Spine J*. 2014 Oct 1;14(10):2299-319.

Daubs MD, Norvell DC, McGuire R, Molinari R, Hermsmeyer JT, Fournay DR, Wolinsky JP, Brodke D. Fusion versus nonoperative care for chronic low back pain: do psychological factors affect outcomes? *Spine (Phila Pa 1976)*. 2011 Oct 1;36(21 Suppl):S96-109.

Foster NE, Mullis R, Hill JC, Lewis M, Whitehurst DGT, Konstantinou, K, Main C, Somerville S, Sowden G, Wathall S, Young J, Hay E. Effect of Stratified Care for Low Back Pain in Family Practice (IMPACT Back): A Prospective Population-Based Sequential Comparison. *Ann Fam Med* 2014; 12(2):102-11.

Kronenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med* 2001; 16(9):606-13.

George SZ, Fritz JM, McNeil DW. Fear-avoidance beliefs as measured by the fear-avoidance beliefs questionnaire: change in fear-avoidance beliefs questionnaire is predictive of change in self-report of disability and pain intensity for patients with acute low back pain. *Clin J Pain* 2006; 22(2):197-203.

Kawchuk GN, Edgecombe TL, Wong AY, Cojocar A, Prasad N; A non-randomized clinical trial to assess the impact of nonrigid, inelastic corsets on spine function in low back pain participants and asymptomatic controls. *Spine J*. 2015 Oct 1;15(10):2222-7.

Morrisette DC, Cholewicki J, Logan S, Seif G, McGowan S; A randomized clinical trial comparing extensible and inextensible lumbosacral orthoses and standard care alone in the management of lower back pain. *Spine (Phila Pa 1976)*. 2014 Oct 1;39(21):1733-42.

van Duijvenbode I, Jellema P, van Poppel M, van Tulder MW. Lumbar supports for prevention and treatment of low back pain. *Cochrane Database of Systematic Reviews* 2008, Issue 2. Art. No.: CD001823.

Chou R, Deyo R, Friedly J, et al. Noninvasive Treatments for Low Back Pain. Comparative Effectiveness Review No. 169. [Internet]. Rockville (MD): Agency for Healthcare Research and Quality (US); 2016 Feb. (Comparative Effectiveness Reviews, No. 169.) [cited 2017 May 4]. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK350276/>

Qaseem A, Wilt TJ, McLean RM, Forciea MA; Clinical Guidelines Committee of the American College of Physicians. Noninvasive Treatments for Acute, Subacute, and Chronic Low Back Pain: A Clinical Practice Guideline From the American College of Physicians. *Ann Intern Med*. 2017;166(7):514-530.

Azadinia F, Ebrahimi E Takamjani, Kamyab M, Parnianpour M, Cholewicki J, Maroufi N. Can lumbosacral orthoses cause trunk muscle weakness? A systematic review of literature. *Spine J*. 2017;17(4):589-602.

About the ABIM Foundation

The mission of the ABIM Foundation is to advance medical professionalism to improve the health care system. We achieve this by collaborating with physicians and physician leaders, medical trainees, health care delivery systems, payers, policymakers, consumer organizations and patients to foster a shared understanding of professionalism and how they can adopt the tenets of professionalism in practice.



To learn more about the ABIM Foundation, visit www.abimfoundation.org.

About the American Chiropractic Association

The American Chiropractic Association (ACA) is the largest professional association in the United States representing doctors of chiropractic. Chiropractors focus on disorders of the musculoskeletal system and the nervous system, and the effects of these disorders on general health and function. Chiropractic services are used most often to treat conditions such as back pain, neck pain, pain in the joints of the arms or legs, and headaches. Widely known for their expertise in spinal manipulation, chiropractors practice a hands-on, drug-free approach to health care that includes patient examination, diagnosis and treatment. On behalf of its members, ACA educates the public about the benefits of chiropractic services, supports research, and provides professional and educational opportunities for chiropractors, with the goal of advancing high-quality patient care. ACA promotes the highest standards of ethics and evidence-informed patient care, and is proud to partner with the *Choosing Wisely*® campaign.



To learn more about ACA, visit www.acatoday.org.