

## Five Things Physicians and Patients Should Question

1

### Do not obtain radiographs in children with bronchiolitis, croup, asthma, or first-time wheezing.

Respiratory illnesses are among the most common reasons for pediatric emergency department (ED) visits, with wheezing being a frequently encountered clinical finding. For children presenting with first-time wheezing or with typical findings of asthma, bronchiolitis, or croup, radiographs rarely yield important positive findings and expose patients to radiation, increased cost of care, and prolonged ED length of stay. National and international guidelines emphasize the value of the history and physical examination in making an accurate diagnosis and excluding serious underlying pathology. Radiography performed in the absence of significant findings has been shown to be associated with overuse of antibiotics. Radiographs should not be routinely obtained in these situations unless findings such as significant hypoxia, focal abnormalities, prolonged course of illness, or severe distress are present. If wheezing is occurring without a clear atopic etiology or without upper respiratory tract infection symptoms (eg, rhinorrhea, nasal congestion, and/or fever), appropriate diagnostic imaging should be considered on a case-by-case basis.

2

### Do not obtain screening laboratory tests in the medical clearance process of pediatric patients who require inpatient psychiatric admission unless clinically indicated.

The incidence of mental health problems in children has increased in the last two decades, with suicide surpassing homicide as the second leading cause of death in teenagers. Most children with acute mental health issues do not have underlying medical etiologies for these symptoms. A large body of evidence, in both adults and children, has shown that routine laboratory testing without clinical indication is unnecessary and adds to health care costs. Any diagnostic testing should be based on a thorough history and physical examination. Universal requirements for routine testing should be abandoned.

3

### Do not order laboratory testing or a CT scan of the head for a patient with an unprovoked, generalized seizure or a simple febrile seizure who has returned to baseline mental status.

Children presenting with unprovoked, generalized seizures or simple febrile seizures who return to their baseline mental status rarely have blood test or CT scan findings that change acute management. CT scans are associated with radiation-related risk of cancer, increased cost of care, and added risk if sedation is required to complete the scan. A head CT scan may be indicated in patients with a new focal seizure, new focal neurologic findings, or high-risk medical history (such as neoplasm, stroke, coagulopathy, sickle cell disease, age <6 months).

4

### Do not obtain abdominal radiographs for suspected constipation.

Functional constipation and nonspecific, generalized abdominal pain are common presenting complaints for children in emergency departments. Constipation is a clinical diagnosis and does not require testing, yet many of these children receive an abdominal radiograph. However, subjectivity and lack of standardization result in poor sensitivity and specificity of abdominal radiographs to diagnose constipation. Use of abdominal radiographs to diagnose constipation has been associated with increased diagnostic error. Clinical guidelines recommend against obtaining routine abdominal radiographs in patients with clinical diagnosis of functional constipation. The diagnosis of constipation or fecal impaction should be made primarily by history and physical examination, augmented by a digital rectal examination when indicated.

5

### Do not obtain comprehensive viral panel testing for patients who have suspected respiratory viral illnesses.

Viral infections occur frequently in children and are a common reason to seek medical care. The diagnosis of a viral illness is made clinically and usually does not require confirmatory testing. Additionally, there is a lack of consistent evidence to demonstrate the impact of comprehensive viral panel (i.e., panels simultaneously testing for 8–20+ viruses) results on clinical outcomes or management, especially in emergency department settings. Hence, most national and international clinical practice guidelines do not recommend their routine use. Additionally, some viral tests are quite expensive, and obtaining nasopharyngeal swab specimens can be uncomfortable for children. Comprehensive viral panel testing can be considered in high-risk patients (eg, immunocompromised) or in situations in which the results will directly influence treatment decisions such as the need for antibiotics, performance of additional tests, or hospitalization. Testing for specific viruses might be indicated if the results of the testing may alter treatment plans (e.g., antivirals for influenza) or public health recommendations (e.g., isolation for SARS-CoV-2). For more specific recommendations related to diagnosis and management of SARS-CoV-2, please see [www.aap.org/en/pages/2019-novel-coronavirus-covid-19-infections/](http://www.aap.org/en/pages/2019-novel-coronavirus-covid-19-infections/).

# How This List Was Created

The American Academy of Pediatrics Section on Emergency Medicine (AAP SOEM) Committee on Quality Transformation (COQT) assembled a task force to oversee the creation of a Pediatric Emergency Medicine *Choosing Wisely* list. The task force first collected suggested recommendations from a diverse group of ED providers (physicians, nurses, and advanced practice providers) from six academic pediatric EDs to gather an initial list of frequently overused and/or avoidable tests and interventions. Task force members independently scored these items on an anchored rating scale based on each item's frequency of overuse in a typical ED shift, the evidence for lack of efficacy, and the potential harm associated with overuse. The scores were discussed, and consensus was reached for the top 25 ranked items. Next, this list of 25 proposed items was sent to all COQT members in a survey format. The COQT member survey respondents selected which 10 items they believed should be included in the *Choosing Wisely* list. The task force then ranked the selected items based on the frequency of selection by COQT members. The five top-ranked items that were not duplicative of items on other subspecialty *Choosing Wisely* lists were submitted and approved by AAP SOEM leadership. The list of five final items with summary evidence was subsequently forwarded for peer review to relevant expert AAP Committee, Council, and Section leadership. The AAP Board of Directors and Executive Committee granted final approval of this list.

## Sources

- Ralston SL, Lieberthal AS, Meissner HC, et al. Clinical practice guideline: the diagnosis, management, and prevention of bronchiolitis. *Pediatrics*. 2014;134(5):e1474-e1502. DOI: <https://doi.org/10.1542/peds.2014-2742>

Trottier ED, Chan K, Allain D, Chauvin-Kimoff L. Managing an acute asthma exacerbation in children. *Paediatr Child Health*. 2021;26(7):438-439. DOI: 10.1093/pch/pxab058

Shah SN, Bachur RG, Simel DL, Neuman MI. Does this child have pneumonia? The rational clinical examination systematic review. *JAMA*. 2017;318(5):462-471. DOI: 10.1001/jama.2017.9039

Schuh S, Lalani A, Allen U, et al. Evaluation of the utility of radiography in acute bronchiolitis. *J Pediatr*. 2007;150(4):429-433. DOI: 10.1016/j.jpeds.2007.01.005

National Heart, Lung, and Blood Institute. Expert Panel Report 4: Guidelines for the Diagnosis and Management of Asthma; National Asthma Education and Prevention Program, Third Expert Panel on the Diagnosis and Management of Asthma. Bethesda, MD: National Heart, Lung, and Blood Institute; 2007:391
- Thrasher TW, Rolli M, Redwood RS, et al. 'Medical clearance' of patients with acute mental health needs in the emergency department: a literature review and practice recommendations. *WMJ*. 2019;118(4):156-163

Donofrio JJ, Horeczko T, Kaji A, Santillanes G, Claudius I. Most routine laboratory testing of pediatric psychiatric patients in the emergency department is not medically necessary. *Health Aff (Millwood)*. 2015;34(5):812-818

Chun TH. Medical clearance: time for this dinosaur to go extinct. *Ann Emerg Med*. 2014;63(6):676-677

Donofrio JJ, Santillanes G, McCammack BD, et al. Clinical utility of screening laboratory tests in pediatric psychiatric patients presenting to the emergency department for medical clearance. *Ann Emerg Med*. 2014;63(6):666-675.e663.

Santillanes G, Donofrio JJ, Lam CN, et al. Is medical clearance necessary for pediatric psychiatric patients? *J Emerg Med*. 2014;46(6):800-807

Santiago LI, Tunik MG, Foltin GL, Mojica MA. Children requiring psychiatric consultation in the pediatric emergency department—epidemiology, resource utilization, and complications. *Pediatr Emerg Care*. 2006;22(2):85-89
- Hirtz D, Ashwal S, Berg A, et al. Practice parameter: Evaluating a first nonfebrile seizure in children. Report of the Quality Standards Subcommittee of the American Academy of Neurology, the Child Neurology Society, and the American Epilepsy Society. *Neurology*. 2000; 55(5):616-623. Reaffirmed October 17, 2020

Rivello JJ Jr, Ashwal S, Hirtz D, et al; American Academy of Neurology Subcommittee; Practice Committee of the Child Neurology Society. Practice parameter: Diagnostic assessment of the child with status epilepticus (an evidence-based review): Report of the Quality Standards Subcommittee of the American Academy of Neurology and the Practice Committee of the Child Neurology Society. *Neurology*. 2006;67(9):1542-1550

McKenzie KC, Hahn CD, Friedman JN; Canadian Paediatric Society, Acute Care Committee. Emergency management of the paediatric patient with convulsive status epilepticus. *Paediatr Child Health*. 2021;26(1):50-57

American Academy of Pediatrics, Subcommittee on Febrile Seizures. Neurodiagnostic evaluation of the children with a simple febrile seizure. *Pediatrics*. 2011;127(2):389-394. DOI: <https://doi.org/10.1542/peds.2010-3318>
- Freedman SB, Rodean J, Hall M, et al. Delayed diagnoses in children with constipation: multicenter retrospective cohort study. *J Pediatr*. 2017;186:87-94.e16. DOI: <https://doi.org/10.1016/j.jpeds.2017.03.061>

Pensabene L, Buonomo C, Fishman L, Chitkara D, Nurko S. Lack of utility of abdominal x-rays in the evaluation of children with constipation: Comparison of different scoring methods. *J Pediatr Gastroenterol Nutr*. 2010;51(2):155-159. DOI: <https://doi.org/10.1097/MPG.0b013e3181cb4309>

Berger MY, Tabbers MM, Kurver MJ, Boluyt N, Benninga MA. Value of abdominal radiography, colonic transit time, and rectal ultrasound scanning in the diagnosis of idiopathic constipation in children: a systematic review. *J Pediatr*. 2012;161(1):44-50.e502. DOI: <https://doi.org/10.1016/j.jpeds.2011.12.045>

Tabbers MM, DiLorenzo C, Berger MY, et al. Evaluation and treatment of functional constipation in infants and children: Evidence-based recommendations from ESPGHAN and NASPGHAN. *J Pediatr Gastroenterol Nutr*. 2014;58(2):258-274. DOI: <https://doi.org/10.1097/mpg.0000000000000266>

Kearney R, Edwards T, Braford M, Klein E. Emergency provider use of plain radiographs in the evaluation of pediatric constipation. *Pediatr Emerg Care*. 2019;35(9):624-629. DOI: 10.1097/PEC.0000000000001549

Freedman SB, Thull-Freedman J, Manson D, et al. Pediatric abdominal radiograph use, constipation, and significant misdiagnoses. *J Pediatr*. 2014;164(1):83-88.e2
- Gill, PJ, Richardson, SE, Ostrow O. Testing for respiratory viruses in children: to swab or not to swab. *JAMA Pediatr*. 2017;171(8):798-804

Noël KC, Fontela PS, Winters N, et al. The clinical utility of respiratory viral testing in hospitalized children: a meta-analysis. *Hosp Pediatr*. 2019;9(7):483-494

Parikh K, Hall M, Mittal V, et al. Establishing benchmarks for the hospitalized care of children with asthma, bronchiolitis, and pneumonia. *Pediatrics*. 2014;134(3):555-562

Innis K, Hasson D, Bodilly L, et al. Do I need proof of the culprit? Decreasing respiratory viral testing in critically ill patients. *Hosp Pediatr*. 2021;11(1):e1-e5

### About the ABIM Foundation

The mission of the ABIM Foundation is to advance medical professionalism to improve the health care system. We achieve this by collaborating with physicians and physician leaders, medical trainees, health care delivery systems, payers, policymakers, consumer organizations and patients to foster a shared understanding of professionalism and how they can adopt the tenets of professionalism in practice.



To learn more about the ABIM Foundation, visit [www.abimfoundation.org](http://www.abimfoundation.org).

### About the American Academy of Pediatrics and the Section on Emergency Medicine

The American Academy of Pediatrics is an organization of 67,000 primary care pediatricians, pediatric medical subspecialists and pediatric surgical specialists dedicated to the health, safety and well-being of infants, children, adolescents and young adults. The AAP Section on Emergency Medicine (SOEM), a group of over 1,900 members, was founded in 1981. The SOEM mission is to sustain, develop, and promote the delivery of optimal emergency care for acutely ill and injured infants, children, and adolescents. The Section envisions: equitable access to emergency care for infants, children, and adolescents; delivery of evidence-guided, safe, and cost-effective emergency care; and development of creative and innovative programs that address the needs of our membership.



### About the Canadian Association of Emergency Physicians (CAEP)

As the national voice of emergency medicine (EM), CAEP provides continuing medical education, advocates on behalf of emergency physicians and their patients, supports research and strengthens the EM community. In co-operation with other specialties and committees, CAEP also plays a vital role in the development of national standards and clinical guidelines. CAEP keeps Canadian emergency physicians informed of developments in the clinical practice of EM and addresses political and societal changes that affect the delivery of emergency health care.



For more information or to see other lists of Things Providers and Patients Should Question, visit [www.choosingwisely.org](http://www.choosingwisely.org).